

Psychometric Analysis of Survey Data


Ron D. Hays, Ph.D.

RCMAR Methods Seminar

February 26, 2024

(3:15-4:15 pm)

<https://uclahs.zoom.us/j/2707654943>



Acknowledgements

- José L. Calderón: “*Mitigating Cognitive Demand for Completion of CAHPS® Surveys by Aging Latinos: CAHPS® Visual Display and Cognitive Design Revision.*” RCMAR Pilot (2015-16 Cohort)
 - National Institute on Aging (Grant Number P30-AG021684)
- Agency for Healthcare and Research Quality (Contract number U18HS029321)
- <https://drjoe.life/obituary/> (November 16, 2023)



AHRQ's CAHPS® Program



- **Agency for Healthcare Research and Quality**
 - ▶ **Research and development agency in the U.S. Department of Health and Human Services**
 - ▶ Since 1995, AHRQ's CAHPS Program has advanced the science of patient experience:
 - Surveys that can be used for high-stakes purposes
 - Quality improvement tools to improve patient experience
 - Research to advance the science of patient experience, including best methods to administer CAHPS surveys and report CAHPS survey findings

The Patient's Voice



CAHPS Surveys reflect the patient's voice.

All CAHPS survey development begins with asking patients/consumers about what's important to measure and report. The resulting survey reflects their input.

CAHPS CG 3.1 Survey

EZ Survey

Form Approved
OMB No. 0935-0124
Exp. Date 1/31/2024

Form Approved
OMB No. 0935-0124
Exp. Date 1/31/2024

Your Experiences with Health Care



Your Health Care

How do you feel about
your health care?

Please let us know!

**Answer this
survey**

This survey is easy to
read and takes about
10 minutes to
complete.

THANK YOU!!



CAHPS Versus “Easy” (EZ) Item

CG-CAHPS 3.1 Item

Passive lead before query ->

In the last 6 months, when you contacted this provider's office to get an appointment for care **you needed right away**, how often did you get an appointment as soon as you needed?

← Random Truncation of Item Lines

Flesh-Kincaid Readability Score:
14th Grade Level
Difficult to Read

EZ Item

How often do you get care as soon as you needed?

← Stanzaic Versification of Item Lines

Flesh-Kincaid Readability Score:
3rd Grade Level
Very Easy to Read

CAHPS Clinician and Group 3.1 Survey

- **31 questions**

- 9 “About You” questions.

- 1 global rating question: Using any number from 0 to 10, where 0 is the worst provider possible, what number would you use to rate this provider?

- Four multi-item scales (12 reports about care items)

Timely Care

6. In the last 6 months, when you contacted this provider’s office to get an appointment for care you needed right away, how often did you get an appointment as soon as you needed?

8. In the last 6 months, when you made an appointment for a check-up or routine care with this provider, how often did you get an appointment as soon as you needed?

10. In the last 6 months, when you contacted this provider’s office during regular office hours, how often did you get an answer to your medical question that same day?

CAHPS Clinician and Group 3.1 Survey

Communication

11. In the last 6 months, how often did this provider explain things in a way that was easy to understand?
12. In the last 6 months, how often did this provider listen carefully to you?
14. In the last 6 months, how often did this provider show respect for what you had to say?
15. In the last 6 months, how often did this provider spend enough time with you?

Coordination of Care

13. In the last 6 months, how often did this provider seem to know the important information about your medical history?

17. In the last 6 months, when this provider ordered a blood test, x-ray, or other test for you, how often did someone from this provider's office follow up to give you those results?

20. In the last 6 months, how often did you and someone from this provider's office talk about all the prescription medicines you were taking?

Office Staff

21. In the last 6 months, how often were clerks and receptionists at this provider's office as helpful as you thought they should be?

22. In the last 6 months, how often did clerks and receptionists at this provider's office treat you with courtesy and respect?

EZ Survey

- **31 questions**

- 9 “About You” questions.

- 1 global rating question: Rate the care this doctor gave you in the last 6 months. Pick a number from 0 to 10. The Worst doctor is 0. The Best doctor is 10

- Four multi-item scales (12 reports about care items)

Timely Care

- 6. How often did you get care as soon as you needed?
- 8. How often did you get an appointment as soon as you needed?
- 10. How often did you get answers to your medical questions the same day?

EZ Survey

- ***Communication***

- 13. How often did this doctor explain things in a way you understood?
- 14. How often did this doctor listen to you carefully?
- 16. How often did this doctor show respect for what you had to say?
- 17. How often did this doctor spend enough time with you?

- ***Coordination of Care***

- 15. How often did this doctor seem to know what is important to you about your health?
- 19. How often did this doctor explain the test results to you?
- 21. How often did this doctor talk about all the medicine you took?

- ***Office Staff***

- 11. How often were clerks and receptionists as helpful as they should be in the last 6 months?
- 12. How often did clerks and receptionists treat you with respect in the last 6 months?

Data Collection

- Safety net healthcare provider in Los Angeles
- CAHPS C-G 3.1 and EZ Paper Surveys
 - Pre-notification letter in advance of survey
 - Personalized letters and survey packets
 - Used first-class postage
 - Sent a second survey to non-respondents
 - English and Spanish surveys
- August 1, 2023, through December 31, 2023.
- n = 264 surveys returned (16 providers)
 - 7 doctors, 5 NPs, 4 PAs

Analysis Plan

- Response rate
- Failure to follow skip patterns
- Missing data
- Item frequencies
- Multi-item scale means (SDs)
- Internal consistency reliability
- Patient-level correlations among measures
- Doctor-level reliability
- Multi-trait scaling
- Factor analyses (exploratory and confirmatory)

Survey response rate

- 18% overall (n = 264)
- 20% for CG 3.1 Survey (n = 147)
- 16% for EZ Survey (n = 117)
- Response rate did not differ significantly by the amount of incentive (\$2 vs \$5)
- Analytic sample (n = 232)
 - n = 133 (CG 3.1) and 99 (EZ) surveys where respondents reported care from sampled provider

Sample Characteristics

| Variable | Percentage |
|-------------------------------|-------------|
| Female | 64% |
| Hispanic | 66% |
| Black | 14% |
| White | 14% |
| Asian | 7% |
| Spanish language survey | 44% |
| High school education or less | 33% |
| Modal age category | 55-64 (40%) |
| Excellent physical health | 15% |
| Very good physical health | 15% |
| Good physical health | 40% |
| Excellent mental health | 24% |
| Very good mental health | 18% |
| Good mental health | 36% |

Percent of Sample Failing to Skip on CG and EZ Survey

| <i>Skip #</i> | <i>CG 3.1 Survey</i> | <i>EZ Survey</i> |
|---------------|----------------------|------------------|
| 1 | 12% (n = 8) | 33% (n = 12) |
| 2 | 38% (n = 8) | 83% (n = 6) |
| 3 | 25% (n = 44) | 39% (n = 28) |
| 4 | 28% (n = 29) | 55% (n = 11) |
| 5 | 25% (n = 55) | 44% (n = 39) |
| 6 | 8% (n = 26) | 55% (n = 11) |
| 7 | 36% (n = 11) | 27% (n = 15) |
| 8 | 5% (n = 132) | 4% (n = 99) |

Item missing data was rare

- 24 items asked of everyone
- Mean missing
 - 0.53 for CG 3.1 survey
 - 1.04 for EZ survey
- $t = 1.78, p = 0.0769$

Percentage of Sample Selecting Each Response Option for CG (EZ) Surveys

| Items | Never | Sometimes | Usually | Always |
|--|---------|-----------|---------|---------|
| <i>Timely Care</i> | | | | |
| Care as soon as needed | 11 (0) | 19 (21) | 25 (37) | 46 (41) |
| Got <u>appointment</u> as soon as needed | 6 (1) | 13 (21) | 36 (40) | 46 (37) |
| Got answers to medical questions <u>same day</u> | 10 (11) | 19 (25) | 21 (22) | 50 (41) |
| <i>Communication</i> | | | | |
| <u>Provider explain</u> things in a way you understand | 2 (3) | 8 (9) | 33 (31) | 57 (57) |
| <u>Provider listen</u> to you carefully | 2 (3) | 6 (7) | 25 (13) | 68 (77) |
| <u>Provider show</u> respect for what you had to say | 3 (1) | 6 (5) | 11 (19) | 80 (75) |
| <u>Provider spend</u> enough time with you | 4 (4) | 5 (6) | 14 (11) | 77 (78) |
| <i>Coordination</i> | | | | |
| <u>Provider know</u> what is important about your health | 2 (2) | 2 (2) | 12 (12) | 83 (83) |
| <u>Provider explain</u> the test results to you | 5 (2) | 6 (9) | 21 (23) | 68 (65) |
| <u>Provider talk</u> about all <u>medicine</u> you take | 4 (2) | 7 (7) | 22 (18) | 67 (73) |
| <i>Office Staff</i> | | | | |
| Clerks and receptionists helpful | 18 (5) | 13 (10) | 12 (13) | 58 (72) |
| Clerks and receptions treat you with respect | 15 (9) | 15 (11) | 25 (11) | 46 (69) |

Patient-Level Means, SDs, Alphas

-----CG-----

-----EZ-----

| | Mean | SD | Alpha* | Mean | SD | Alpha* |
|---------------|------|------|--------|------|------|--------|
| Timely | 3.09 | 0.91 | 0.90 | 3.06 | 0.83 | 0.81 |
| Communication | 3.65 | 0.65 | 0.92 | 3.65 | 0.62 | 0.83 |
| Coordination | 3.22 | 0.80 | 0.65 | 3.52 | 0.75 | 0.80 |
| Office Staff | 3.52 | 0.65 | 0.83 | 3.53 | 0.73 | 0.88 |

* $(MS_{BMS} - MS_{EMS}) / MS_{BMS}$ MS_{EMS} = Patient x Item interaction

Correlations Among Scales and Global Doctor Rating (CG above and EZ below diagonal)

| | <i>Timely</i> | <i>Communication</i> | <i>Coordination</i> | <i>Office Staff</i> | <i>Global Rating</i> |
|----------------------|---------------|----------------------|---------------------|---------------------|----------------------|
| <i>Timely</i> | 1.00 | 0.26 | 0.36 | 0.28 | 0.31 |
| <i>Communication</i> | 0.34 | 1.00 | 0.61 | 0.48 | 0.73 |
| <i>Coordination</i> | 0.40 | 0.82 | 1.00 | 0.48 | 0.46 |
| <i>Office Staff</i> | 0.38 | 0.53 | 0.53 | 1.00 | 0.42 |
| <i>Global Rating</i> | 0.37 | 0.73 | 0.77 | 0.33 | 1.00 |

Provider-Level Reliability Estimates

----- CG ----- ----- EZ ----- -----

| | Reliability | N for 0.70 | Reliability | N for 0.70 |
|----------------------|-------------|------------|-------------|------------|
| <i>Timely</i> | 0.00 | ∞ | 0.44 | 19 |
| <i>Communication</i> | 0.00 | ∞ | 0.47 | 17 |
| <i>Coordination</i> | 0.08 | 207 | 0.19 | 65 |
| <i>Office Staff</i> | 0.00 | ∞ | 0.00 | ∞ |
| <i>Global Rating</i> | 0.38 | 31 | 0.41 | 21 |

$(MS_{BMS} - MS_{WMS}) / MS_{BMS}$ Where MS_{WMS} = Within mean square.

MULTI -- MULTITRAIT SCALING PROGRAM

CAHPS CG SURVEY

| item | Timely3 | Comm4 | Coord3 | Couresp2 |
|------|---------|-------|--------|----------|
| Q6 | 0.82* | 0.30 | 0.32 | 0.35 |
| Q8 | 0.84* | 0.26 | 0.37 | 0.39 |
| Q10 | 0.71* | 0.25 | 0.31 | 0.39 |
| Q11 | 0.26 | 0.88* | 0.59 | 0.53 |
| Q12 | 0.26 | 0.88* | 0.61 | 0.54 |
| Q14 | 0.23 | 0.76* | 0.49 | 0.42 |
| Q15 | 0.28 | 0.75* | 0.46 | 0.39 |
| Q13 | 0.30 | 0.82 | 0.42* | 0.50 |
| Q17 | 0.29 | 0.35 | 0.59* | 0.40 |
| Q20 | 0.27 | 0.35 | 0.57* | 0.36 |
| Q21 | 0.39 | 0.42 | 0.50 | 0.70* |
| Q22 | 0.34 | 0.54 | 0.46 | 0.70* |

How often did this provider seem to know the important information about medical history?

MULTI -- MULTITRAIT SCALING PROGRAM

CAHPS CG SURVEY

Primary Care First Patient Experience of Care Survey:
<https://pcfpecs.org/General-Information/About-PCF-PECS>

| item | Timely3 | Comm4 | Coord3 | Couresp2 |
|------|---------|-------|--------|----------|
| Q6 | 0.82* | 0.30 | 0.32 | 0.35 |
| Q8 | 0.84* | 0.26 | 0.37 | 0.39 |
| Q10 | 0.71* | 0.25 | 0.31 | 0.39 |
| Q11 | 0.26 | 0.88* | 0.59 | 0.53 |
| Q12 | 0.26 | 0.88* | 0.61 | 0.54 |
| Q14 | 0.23 | 0.76* | 0.49 | 0.42 |
| Q15 | 0.28 | 0.75* | 0.46 | 0.39 |
| Q13 | 0.30 | 0.82 | 0.42* | 0.50 |
| Q17 | 0.29 | 0.35 | 0.59* | 0.40 |
| Q20 | 0.27 | 0.35 | 0.57* | 0.36 |
| Q21 | 0.39 | 0.42 | 0.50 | 0.70* |
| Q22 | 0.34 | 0.54 | 0.46 | 0.70* |

How often did this provider seem to know the important information about medical history?

Exploratory Factor Analysis

CG Survey
PRINCIPAL COMPONENTS ANALYSIS
GUTTMAN'S WEAKEST LOWER BOUND
LOOKING FOR EIGENVALUES ≥ 1

The FACTOR Procedure
Initial Factor Method: Principal Components

Prior Communality Estimates: ONE

| Eigenvalues of the Correlation Matrix: Total = 12 Average = 1 | | | | |
|--|------------|------------|------------|------------|
| | Eigenvalue | Difference | Proportion | Cumulative |
| 1 | 5.61993719 | 3.44114111 | 0.4683 | 0.4683 |
| 2 | 2.17879608 | 1.12350641 | 0.1816 | 0.6499 |
| 3 | 1.05528967 | 0.12358322 | 0.0879 | 0.7378 |
| 4 | 0.93170645 | 0.40698858 | 0.0776 | 0.8155 |
| 5 | 0.52471787 | 0.09175144 | 0.0437 | 0.8592 |
| 6 | 0.43296642 | 0.04691628 | 0.0361 | 0.8953 |
| 7 | 0.38605015 | 0.07254840 | 0.0322 | 0.9275 |
| 8 | 0.31350174 | 0.06237872 | 0.0261 | 0.9536 |
| 9 | 0.25112302 | 0.11509493 | 0.0209 | 0.9745 |
| 10 | 0.13602809 | 0.01477922 | 0.0113 | 0.9858 |
| 11 | 0.12124888 | 0.07261443 | 0.0101 | 0.9959 |
| 12 | 0.04863445 | | 0.0041 | 1.0000 |

7.31, 1.97, and 0.98
eigenvalues for polychoric
correlations

PARALLEL.EXE: LATENT ROOTS OF RANDOM DATA CORRELATION MATRICES PROGRAM
PROGRAMMER: RON HAYS, RAND CORPORATION
FOR 131 SUBJECTS AND 12 VARIABLES AND 100 SAMPLES

Hays, R. D. (1987). PARALLEL: A program for performing parallel
analysis. Applied Psychological Measurement, 11, 58.

EIGENVALUES FOR FACTOR ANALYSIS SMC ESTIMATES FOLLOW:

| | OBSERVED | RANDOM | SLOPE | |
|--------|-------------|----------|-----------|-----|
| | ===== | ===== | ===== | |
| LAMBDA | 1= 5.129700 | 0.668903 | ----- | |
| | | | -3.028500 | |
| LAMBDA | 2= 2.101200 | 0.489875 | ----- | |
| | | | -1.416800 | *** |
| LAMBDA | 3= 0.684400 | 0.393931 | ----- | |
| | | | -0.185500 | |
| LAMBDA | 4= 0.498900 | 0.323764 | ----- | |
| | | | -0.287800 | *** |
| LAMBDA | 5= 0.211100 | 0.234022 | ----- | |
| | | | -0.034300 | |
| LAMBDA | 6= 0.176800 | 0.157072 | ----- | |


(CAN'T COMPUTE LAMBDA 7 :LOG OF ZERO OR NEGATIVE IS UNDEFINED)

Results of Parallel Analysis Indicate Maximum of 4 Factors.
Slopes followed by asterisks indicate discontinuity points
that may be suggestive of the number of factors to retain.

Scree Plot


The FACTOR Procedure
Initial Factor Method: Principal Factors

Scree Plot of Eigenvalues



Tucker/Lewis reliability
Coefficient for 3 factors = 0.80
and 4 factors = 0.82

Polychoric Correlations



**OBLIQUE PROMAX ROTATION -- 4 FACTOR SOLUTION satisfaction sco
COMMON FACTOR ANALYSIS
SQUARED MULTIPLE CORRELATIONS AS PRIOR COMMUNALITIES ES**

**The FACTOR Procedure
Rotation Method: Promax (power = 3)**

| Rotated Factor Pattern (Standardized Regression Coefficients) | | | | | |
|---|-----|----------|----------|----------|----------|
| | | Factor1 | Factor2 | Factor3 | Factor4 |
| Q11 | Q11 | 0.90162 | -0.02213 | 0.07187 | -0.01761 |
| Q12 | Q12 | 0.87963 | -0.05304 | 0.07360 | 0.07399 |
| Q13 | Q13 | 0.83030 | 0.02288 | 0.02859 | 0.04306 |
| Q14 | Q14 | 0.81941 | 0.00383 | 0.02628 | -0.06644 |
| Q15 | Q15 | 0.81097 | 0.04420 | -0.11529 | 0.07241 |
| Q8 | Q8 | -0.03014 | 0.93497 | -0.02639 | 0.05938 |
| Q6 | Q6 | 0.15636 | 0.92633 | 0.00326 | -0.15567 |
| Q10 | Q10 | -0.14797 | 0.60869 | 0.05063 | 0.23116 |
| Q22 | Q22 | 0.10378 | -0.01726 | 0.78082 | -0.05110 |
| Q21 | Q21 | 0.01794 | 0.05482 | 0.74874 | 0.06879 |
| Q20 | Q20 | 0.09261 | -0.01493 | -0.06442 | 0.71117 |
| Q17 | Q17 | 0.02310 | 0.13730 | 0.12152 | 0.60273 |

| Inter-Factor Correlations | | | | |
|---------------------------|---------|---------|---------|---------|
| | Factor1 | Factor2 | Factor3 | Factor4 |
| Factor1 | 1.00000 | 0.27320 | 0.56104 | 0.36463 |
| Factor2 | 0.27320 | 1.00000 | 0.41458 | 0.44485 |
| Factor3 | 0.56104 | 0.41458 | 1.00000 | 0.49657 |
| Factor4 | 0.36463 | 0.44485 | 0.49657 | 1.00000 |

**OBLIQUE PROMAX ROTATION -- 4 FACTOR SOLUTION
COMMON FACTOR ANALYSIS
SQUARED MULTIPLE CORRELATIONS AS PRIOR COMMUNALITIES ESTIMATES**

The FACTOR Procedure
Rotation Method: Promax (power = 3)

| Rotated Factor Pattern (Standardized Regression Coefficients) | | | | | |
|---|-----|----------|----------|----------|----------|
| | | Factor1 | Factor2 | Factor3 | Factor4 |
| Q15 | Q15 | 0.97694 | 0.09429 | -0.04157 | -0.23714 |
| Q13 | Q13 | 0.93432 | 0.08869 | -0.00913 | -0.01028 |
| Q14 | Q14 | 0.93396 | -0.08749 | -0.00360 | 0.06967 |
| Q12 | Q12 | 0.91413 | 0.00053 | 0.07142 | 0.06640 |
| Q11 | Q11 | 0.89476 | 0.04113 | 0.12845 | -0.01225 |
| Q20 | Q20 | 0.89372 | -0.07638 | 0.02099 | 0.25091 |
| Q6 | Q6 | 0.06504 | 0.99723 | -0.04723 | -0.03030 |
| Q8 | Q8 | 0.00382 | 0.91485 | 0.02135 | 0.10993 |
| Q21 | Q21 | 0.00885 | -0.00072 | 0.94597 | 0.02325 |
| Q22 | Q22 | 0.19383 | -0.02382 | 0.87073 | -0.03926 |
| Q17 | Q17 | 0.11339 | -0.02798 | -0.10549 | 0.95409 |
| Q10 | Q10 | -0.16854 | 0.25073 | 0.22476 | 0.64823 |

Standardized Factor Loadings for 3-Factor Categorical Factor Analytic Model

| Item | Timely | Communication | Office |
|--|--------|---------------|--------|
| Q6 Care as soon as needed | 0.92 | | |
| Q8 Got an appointment as soon as needed | 0.95 | | |
| Q10 Got answers to questions <u>same day</u> | 0.75 | | |
| Q11 Explained things in a way you understand | | 0.94 | |
| Q12 Listen to you carefully | | 0.94 | |
| Q14 Show respect for what you had to say | | 0.80 | |
| Q15 Spent enough time with you | | 0.77 | |
| Q13 Know what is important about your health | | 0.87 | |
| Q17 Explain the test results to you | | 0.32 | |
| Q20 Talk about the medicine you take | | 0.56 | |
| Q21 Staff helpful | | | 0.82 |
| Q22 Staff courtesy and respect | | | 0.88 |

Comparative fit index = 0.95; RMSEA = 0.09





FIG. 1. Item characteristic curves for CAHPS Communication Items.

Email: drhays@ucla.edu

<https://labs.dgsom.ucla.edu/hays/pages/>



Cappelleri, J. C., Lundy, J.J., & Hays, R. D. (2014). Overview of classical test theory and item response theory for quantitative assessment of items in developing patient-reported outcome measures. Clinical Therapeutics., 36 (5), 648-662.

Hays, R. D., Walling, A. M., Sudore, R. L., Chau, A., & Wenger, N. S. (2023). Support for the use of Consumer Assessment of Healthcare Providers and Systems communication items among seriously ill patients. J Palliat Med, 26(9), 1234-1239.

Appendix: Confirmatory Factor Analysis Fit Indices

- Normed fit index: $\frac{\chi_{null}^2 - \chi_{model}^2}{\chi_{null}^2}$
 - Non-normed fit index: $\frac{\frac{\chi_{null}^2}{df_{null}} - \frac{\chi_{model}^2}{df_{model}}}{\left[\frac{\chi_{null}^2}{df_{null}} - 1 \right]}$
 - Comparative fit index: $1 - \left[\frac{\chi_{model}^2 - df_{model}}{\chi_{null}^2 - df_{null}} \right]$
- RMSEA = SQRT $(\lambda^2 - df)/SQRT (df (N - 1))$

CFI ≥ 0.95 and RMSEA ≤ 0.06